

WEDGE MODULES FOR TWO-PARAMETER QUANTUM GROUPS

NAIHUAN JING, LILI ZHANG, MING LIU*

ABSTRACT. The Yang-Baxterization $R(z)$ of the trigonometric R-matrix is computed for the two-parameter quantum affine algebra $U_{r,s}(\widehat{\mathfrak{sl}}_n)$. Using the fusion procedure we construct all fundamental representations of the quantum algebra $U_{r,s}(\mathfrak{sl}_n)$ as (r, s) -wedge products of the natural representation.

1. INTRODUCTION

Two-parameter general linear and special linear quantum groups were introduced by Takeuchi [15] in 1990. This new type of quantum groups was used in [4] to incorporate two seemingly different types of quantum general linear groups: the usual quantum general linear groups from Drinfeld-Jimbo quantum algebras [6, 9] and Dipper-Donkin quantum linear groups [5] connected with quantum q -Schur algebras. Similar two-parametric quantum groups originated from special solutions of Yang-Baxter equations was also studied in [10] and they interpolate the usual quantum groups and the exotic cases from the six vertex model. All these quantum groups can be viewed as quantum transformation groups over certain quantum planes in the sense of Manin [13].

In 2001 Benkart and Witherspoon [1] investigated the two-parameter quantum group in connection with the down-up algebras. In [2] Benkart and Witherspoon developed the two-parameter quantum groups corresponding to general linear and special linear Lie algebras $\mathfrak{gl}_n$ and $\mathfrak{sl}_n$, and constructed the corresponding R -matrix and the quantum Casimir element. They further showed that these two algebras can be realized as Drinfeld doubles. In [3] the representation theory of two-parameter quantum groups $U_{r,s}(\mathfrak{gl}_n)$ and $U_{r,s}(\mathfrak{sl}_n)$ was studied and an explicit description of the (r,s) -symmetric tensor $S_{r,s}^2(V)$ and the R -matrix $R = R_{VV}$ was given.

The fundamental representations play an important role in the representation theory of classical Lie algebras serving as building blocks and examples for general highest weight representations. In the case of $\mathfrak{sl}_n$, the fundamental representations are just the wedge modules of the natural representation. The fundamental representations of the quantum group $U_q(\mathfrak{sl}_n)$ were first studied by Rosso in [14] using the general

MSC (2010): Primary: 17B30; Secondary: 17B68.

Keywords: two-parameter quantum groups, Yang-Baxterization, (r,s) -wedge module.

*Corresponding author.

highest weight theory. Later the fundamental modules were reconstructed using the fusion procedure [12, 11] in connection with the quantum affine algebras $U_q(\widehat{\mathfrak{sl}}_n)$. A natural question of the wedge modules arises for the two-parameter quantum groups. Although the tensor modules of $U_{r,s}(\widehat{\mathfrak{sl}}_n)$ have been constructed in [3] it is nontrivial to pass down to the irreducible quotient modules, and the best method seems to be the fusion procedure. In order to carry out the fusion procedure one first needs to find the R -matrix with spectral parameter as in the one-parameter case.

In this paper we first use the Yang-Baxterization method developed by Ge-Wu-Xue [7] to construct a spectral parameter dependent R -matrix for the two-parameter quantum algebra $U_{r,s}(\widehat{\mathfrak{sl}}_n)$ from Benkart-Witherspoon R -matrix. This is done by exploiting the two-parameter R -matrix as a braid group representation. We remark that our R -matrix $R(z)$ can also be viewed as the R -matrix of two-parameter quantum affine algebra $U_{r,s}(\widehat{\mathfrak{sl}}_n)$ [8], from which one can use the fusion procedure to study the (r,s) -wedge modules of $U_{r,s}(\mathfrak{sl}_n)$.

The paper is organized as follows. In section 2, we give a brief introduction of two-parameter quantum groups $U_{r,s}(\mathfrak{gl}_n)$ and $U_{r,s}(\mathfrak{sl}_n)$ and recall the results given in [3]. In section 3, we obtain an R -matrix with spectral parameter which can be regarded as the R -matrix corresponding to the two-parameter quantum affine algebra $U_{r,s}(\widehat{\mathfrak{sl}}_n)$ by using the Yang-Baxterization method of Ge, Wu and Xue. In section 4 we determine all (r,s) -wedge modules of $U_{r,s}(\mathfrak{sl}_n)$.

2. TWO-PARAMETER QUANTUM GROUP $U_{r,s}(\mathfrak{sl}_n)$ AND R -MATRIX

We first recall the definition of the two-parameter quantum group $U_{r,s}(\mathfrak{sl}_n)$ and some basics about their representations from [3]. Let $\epsilon_1, \epsilon_2, \dots, \epsilon_n$ denote an orthonormal basis of a Euclidean space E with an inner product $\langle \cdot, \cdot \rangle$. Let $\Pi = \{\alpha_j = \epsilon_j - \epsilon_{j+1} | j = 1, 2, \dots, n-1\}$ be the set of the simple roots of type A_{n-1} , then $\Phi = \{\epsilon_i - \epsilon_j | 1 \leq i \neq j \leq n\}$ is the set of all roots.

Fix nonzero elements $r, s \in \mathbb{C}$ with $r \neq s$.

Definition 2.1. The two-parameter quantized enveloping algebra $U_{r,s}(\mathfrak{sl}_n)$ is the unital associative algebra over $\mathbb{C}$ generated by $e_i, f_i, \omega_i, \omega'_i$, $1 \leq i < n$ with the following relations:

- (R1). The generators ω_i, ω'_i are invertible commutative elements,
- (R2). $\omega_i e_j = r^{\langle \epsilon_i, \alpha_j \rangle} s^{\langle \epsilon_{i+1}, \alpha_j \rangle} e_j \omega_i$ and $\omega_i f_j = r^{-\langle \epsilon_i, \alpha_j \rangle} s^{-\langle \epsilon_{i+1}, \alpha_j \rangle} f_j \omega_i$,
- (R3). $\omega'_i e_j = r^{\langle \epsilon_{i+1}, \alpha_j \rangle} s^{\langle \epsilon_i, \alpha_j \rangle} e_j \omega'_i$ and $\omega'_i f_j = r^{-\langle \epsilon_{i+1}, \alpha_j \rangle} s^{-\langle \epsilon_i, \alpha_j \rangle} f_j \omega'_i$,
- (R4). $[e_i, f_j] = \frac{\delta_{ij}}{r-s}(\omega_i - \omega'_i)$,
- (R5). $[e_i, e_j] = [f_i, f_j] = 0, i f | i - j | > 1$,
- (R6). $e_i^2 e_{i+1} - (r+s) e_i e_{i+1} e_i + r s e_{i+1} e_i^2 = 0$ and $e_{i+1}^2 e_i - (r+s) e_{i+1} e_i e_{i+1} + r s e_{i+1}^2 e_i = 0$,
- (R7). $f_i^2 f_{i+1} - (r^{-1} + s^{-1}) f_i f_{i+1} f_i + r^{-1} s^{-1} f_{i+1} f_i^2 = 0$ and $f_{i+1}^2 f_i - (r^{-1} + s^{-1}) f_{i+1} f_i f_{i+1} + r^{-1} s^{-1} f_{i+1}^2 f_i = 0$, where $[\cdot, \cdot]$ is the usual commutator.

Remark 2.2. When $r = q$, $s = q^{-1}$, the algebra modulo the ideal generated by the elements $\omega_j - \omega'_j$, $1 \leq j < n$, is $U_q(\mathfrak{sl}_n)$.

The algebra $U_{r,s}(\mathfrak{sl}_n)$ is a Hopf algebra under the coproduct that ω_i, ω'_i are group-like elements and other nontrivial coproducts, counits and antipodes are given by:

$$\Delta(e_i) = e_i \otimes 1 + \omega_i \otimes e_i, \Delta(f_i) = 1 \otimes f_i + f_i \otimes \omega'_i,$$

$$\epsilon(e_i) = \epsilon(f_i) = 0, S(e_i) = -\omega_i^{-1}e_i, S(f_i) = -f_i\omega'_i{}^{-1}.$$

The natural representation of $U = U_{r,s}(\mathfrak{sl}_n)$ can be easily described as follows. Let $\Lambda = \mathbb{Z}\epsilon_1 \oplus \mathbb{Z}\epsilon_2 \oplus \cdots \oplus \mathbb{Z}\epsilon_n$ be the weight lattice of $\mathfrak{gl}_n$ and $Q = \mathbb{Z}\Phi$ the root lattice. Assume Λ is equipped with a partial order in which $\nu \leq \lambda$ if and only if $\lambda - \nu \in \sum_{i=1}^{n-1} \mathbb{Z}_{\geq 0}\alpha_i$. For each $i < n$ define the fundamental weight

$$\bar{\omega}_i = \epsilon_1 + \cdots + \epsilon_i.$$

Then $\Lambda_{\mathfrak{sl}} = \mathbb{Z}\bar{\omega}_1 \oplus \cdots \oplus \mathbb{Z}\bar{\omega}_{n-1}$ is the weight lattice of $\mathfrak{sl}$.

Corresponding to $\lambda \in \Lambda_{\mathfrak{sl}}$, there is an algebra homomorphism $\hat{\lambda}$ from the subalgebra U_0 of $U_{r,s}(\mathfrak{sl}_n)$ generated by $\omega_i^{\pm 1}, \omega'_i{}^{\pm 1}$ to $\mathbb{C}$ given by:

$$(2.1) \quad \hat{\lambda}(\omega_i) = r^{\langle \epsilon_i, \lambda \rangle} s^{\langle \epsilon_{i+1}, \lambda \rangle}, \quad \hat{\lambda}(\omega'_i) = r^{\langle \epsilon_{i+1}, \lambda \rangle} s^{\langle \epsilon_i, \lambda \rangle}.$$

Let V be the n -dimensional vector space over $\mathbb{C}$ with basis $\{v_j | 1 \leq j \leq n\}$. The natural representation is the $U_{r,s}(\mathfrak{sl}_n)$ -module V with the action given by:

$$\begin{aligned} e_j &= E_{j,j+1}, \quad f_j = E_{j+1,j}, \\ \omega_j &= rE_{jj} + sE_{j+1,j+1} + \sum_{k \neq j, j+1} E_{kk}, \\ \omega'_j &= sE_{jj} + rE_{j+1,j+1} + \sum_{k \neq j, j+1} E_{kk}, \end{aligned}$$

where $1 \leq j \leq n-1$.

It is clear that

$$(2.2) \quad \omega_i v_j = r^{\langle \epsilon_i, \epsilon_j \rangle} s^{\langle \epsilon_{i+1}, \epsilon_j \rangle},$$

$$(2.3) \quad \omega'_i v_j = r^{\langle \epsilon_{i+1}, \epsilon_j \rangle} s^{\langle \epsilon_i, \epsilon_j \rangle},$$

for all i, j , so v_j has weight $\epsilon_j = \bar{\omega}_1 - (\alpha_1 + \cdots + \alpha_{j-1})$. Therefore $V = \bigoplus_{j=1}^n V_{\epsilon_j}$ is the weight decomposition, and it is a simple module of $U_{r,s}(\mathfrak{sl}_n)$.

In [3] the authors studied the tensor powers of the natural representation V of $U_{r,s}(\mathfrak{sl}_n)$ and the associated R -matrix. Let $R = R_{VV}$ be the R -matrix associated to V given by the coproduct, and for $1 \leq i < k$, let R_i be the $U_{r,s}(\mathfrak{sl}_n)$ module isomorphism on $V^{\otimes k}$ defined by

$$R_i(w_1 \otimes w_2 \otimes \cdots \otimes w_k) = w_1 \otimes \cdots \otimes R(w_i \otimes w_{i+1}) \otimes w_{i+2} \otimes \cdots \otimes w_k.$$

Since $R = R_{VV}$ satisfies the Yang-Baxter equation, the braid relations hold:

$$R_i R_{i+1} R_i = R_{i+1} R_i R_{i+1} \text{ for } 1 \leq i < k,$$

The construction also implies that for $|i - j| \geq 2$,

$$R_i R_j = R_j R_i.$$

The following result was given in [3].

Proposition 2.3. *Whenever $s \neq -r$, the $U_{r,s}(\mathfrak{sl}_n)$ -module $V \otimes V$ decomposes into two simple submodules, $S_{r,s}^2(V)$ (the (r,s) -symmetric tensors), and $\Lambda_{r,s}^2(V)$ (the (r,s) -antisymmetric tensors). These modules are defined as follows:*

- (i): $S_{r,s}^2(V)$ is the span of $\{v_i \otimes v_i | 1 \leq i \leq n\} \cup \{v_i \otimes v_j + sv_j \otimes v_i | 1 \leq i < j \leq n\}$.
- (ii): $\Lambda_{r,s}^2(V)$ is the span of $\{v_i \otimes v_j - rv_j \otimes v_i | 1 \leq i < j \leq n\}$.

Consequently one has that

$$(2.4) \quad R_i^2 = (1 - rs^{-1})R_i + rs^{-1}Id$$

for all $1 \leq i < k$.

Proposition 2.4. [3] *The minimum polynomial of R on $V \otimes V$ is $(t - 1)(t + rs^{-1})$ if $s \neq -r$.*

From the proposition 2.4, it follows that the action of R on $V \otimes V$ is given as follows:

$$(2.5) \quad R = \sum_{i=1}^n E_{ii} \otimes E_{ii} + r \sum_{i < j} E_{ji} \otimes E_{ij} + s^{-1} \sum_{i < j} E_{ij} \otimes E_{ji} + (1 - rs^{-1}) \sum_{i < j} E_{jj} \otimes E_{ii}.$$

3. YANG-BAXTERIZATION

Starting from a R -matrix of the Yang-Baxter equation (YBE), Yang-Baxterization of R recovers the associated spectral parameter dependent R -matrix $R(z)$ satisfying the YBE:

$$R_1(z)R_2(zw)R_1(w) = R_2(w)R_1(zw)R_2(z).$$

When R has two or three eigenvalues Ge, Wu and Xue [7] gave an algorithm to compute the Yang-Baxterization of R . Suppose R has two distinct eigenvalue λ_1, λ_2 , the Yang-Baxterization gives the resulting R -matrix $R(z)$ as follows:

$$(3.1) \quad R(z) = \lambda_2^{-1}R + z\lambda_1 R^{-1}.$$

Proposition 2.4 shows that $R = R_{VV}$ has eigenvalues 1 and $-rs^{-1}$ on $V \otimes V$, then by the Yang-Baxterization we have the following theorem. We remark that the resulting R -matrix with spectral parameter z is (r, s) -analogue of the R -matrix for the quantum affine algebra $U_q(\widehat{\mathfrak{sl}}_n)$.

Theorem 3.1. *For the braid group representation $R = R_{VV}$, the resulting R -matrix $R(z)$ is given as follows:*

$$(3.2) \quad \begin{aligned} R(z) = & (1 - zrs^{-1}) \sum_{i=1}^n E_{ii} \otimes E_{ii} + (1 - z) \left(r \sum_{i>j} + s^{-1} \sum_{i<j} \right) E_{ij} \otimes E_{ji} \\ & + z(1 - rs^{-1}) \sum_{i<j} E_{ii} \otimes E_{jj} + (1 - rs^{-1}) \sum_{i>j} E_{ii} \otimes E_{jj}. \end{aligned}$$

Proof. We already mentioned that $R = R_{VV}$ has two distinct eigenvalues $\lambda_1 = -rs^{-1}$ and $\lambda_2 = 1$. Ge-Wu-Xue's Yang-Baxterization implies that the resulting R -matrix with spectral parameter is then given by:

$$R(z) = R - zrs^{-1}R^{-1}$$

On the other hand, Eq. (2.4) implies that

$$R^{-1} = r^{-1}sR + (1 - r^{-1}s)I.$$

Thus $R(z) = (1 - z)R - z(rs^{-1} - 1)I$, which is exactly Eq. (3.2). $\square$

Remark 3.2. Clearly $R(0) = R$. Moreover, when $r = q$ and $s = q^{-1}$, the R -matrix $R(z)$ turns into

$$R_q(z) = (1 - zq^2) \sum_{i=1}^n E_{ii} \otimes E_{ii} + (1 - z)q \sum_{i \neq j} E_{ij} \otimes E_{ji} + (1 - q^2) \left(\sum_{i>j} + z \sum_{i<j} \right) E_{ii} \otimes E_{jj},$$

which is exactly the R -matrix for the quantum affine algebra $U_q(\mathfrak{sl}_n)$. In this regard we can view $R(z)$ as an (r, s) -analogue of the R -matrix $R_q(z)$ of the quantum affine algebra $U_q(\mathfrak{sl}_n)$ [8].

4. (r, s) -WEDGE MODULES OF $U_{r,s}(sl_n)$

In [3] Benkart and Witherspoon studied the (r, s) -symmetric tensor $S_{r,s}^2(V)$ (see Proposition 2.3) using the Hopf algebra structure. Here we give alternative description of the symmetric tensors using the fusion procedure.

Proposition 4.1. *The subspace $S_{r,s}^2(V)$ is equal to the image of $R(rs^{-1})$ on $V \otimes V$, and $\Lambda_{r,s}^2(V)$ is equal to the kernel of $R(r^{-1}s)$ on $V \otimes V$.*

Proof. This is a simple consequence of the spectral decomposition of the R -matrix $R(z)$. In fact applying $R(rs^{-1})$ to $v_i \otimes v_j$ proves the first statement and the second one is verified by evaluating $R(r^{-1}s)(v_i \otimes v_j - rv_j \otimes v_i) = 0$ for $i < j$. $\square$

Remark 4.2. The above proposition is the analogue of the one-parameter case (see [11]). In fact, when $s^{-1} = q = r$, the (r, s) -symmetric tensor $S_{r,s}^2(V)$ of $U_{r,s}(\mathfrak{sl}_n)$ reduces to the q -symmetric tensor W of $U_q(\mathfrak{sl}_n)$ (see [11]).

Now we can study the general wedge products of the two-parameter case using the fusion procedure as in [11]. The following theorem completely determine (r, s) -wedge modules for $U_{r,s}(\mathfrak{sl}_n)$.

Theorem 4.3. *On the tensor product of the natural representation of $U_{r,s}(\mathfrak{sl}_n)$ we have*

$$(4.1) \quad V^{\otimes k} / \sum_{i=0}^{k-2} V^{\otimes i} \otimes S_{r,s}^2(V) \otimes V^{\otimes(k-i-2)} \cong V(\overline{\omega}_k)$$

Proof. We denote the image of $v_{i_1} \otimes v_{i_2} \cdots \otimes v_{i_k} \in V^{\otimes k}$ in $V^{\otimes k} / \sum_{i=0}^{k-2} V^{\otimes i} \otimes S_{r,s}^2(V) \otimes V^{\otimes(k-i-2)}$ by $v_{i_1} \wedge v_{i_2} \wedge \cdots \wedge v_{i_k}$.

First we show that the vector $v_{i_1} \wedge v_{i_2} \wedge \cdots \wedge v_{i_k}$ is a highest vector with highest weight $\overline{\omega}_k$ in the $U_{r,s}(\mathfrak{sl}_n)$ module.

It follows from Eqs. (2.2)–(2.3) and the Hopf algebra structure of $U_{r,s}(\mathfrak{sl}_n)$ that

$$\begin{aligned} \omega_i \cdot v_1 \wedge v_2 \wedge \cdots \wedge v_k &= r^{<\varepsilon_i, \varepsilon_1 + \varepsilon_2 + \cdots + \varepsilon_k>} s^{<\varepsilon_{i+1}, \varepsilon_1 + \varepsilon_2 + \cdots + \varepsilon_k>} v_1 \wedge v_2 \wedge \cdots \wedge v_k, \\ \omega'_i \cdot v_1 \wedge v_2 \wedge \cdots \wedge v_k &= r^{<\varepsilon_{i+1}, \varepsilon_1 + \varepsilon_2 + \cdots + \varepsilon_k>} s^{<\varepsilon_i, \varepsilon_1 + \varepsilon_2 + \cdots + \varepsilon_k>} v_1 \wedge v_2 \wedge \cdots \wedge v_k. \end{aligned}$$

Hence $v_1 \wedge v_2 \wedge \cdots \wedge v_k$ has weight $\overline{\omega}_k = \varepsilon_1 + \varepsilon_2 + \cdots + \varepsilon_k$.

Furthermore the action of e_i on $V^{\otimes k}$ is given by

$$\begin{aligned} e_i \cdot v_1 \otimes v_2 \otimes \cdots \otimes v_k &= \sum_{j=1}^k \omega_i v_1 \otimes \cdots \otimes \omega_i v_{j-1} \otimes e_i v_j \otimes v_{j+1} \otimes \cdots \otimes v_k \\ &= \sum_{j=1}^k r^{<\varepsilon_i, \varepsilon_1 + \cdots + \varepsilon_{j-1}>} s^{<\varepsilon_{i+1}, \varepsilon_1 + \cdots + \varepsilon_{j-1}>} \delta_{j,i+1} v_1 \cdots \otimes v_{j-1} \otimes v_i \otimes v_{j+1} \cdots \otimes v_k \\ &= r v_1 \otimes \cdots \otimes v_i \otimes v_i \otimes v_{i+2} \otimes v_{i+3} \otimes \cdots \otimes v_k \quad \text{or} \quad 0 \\ &\in \sum_{j=0}^{k-2} V^{\otimes j} \otimes S_{r,s}^2(V) \otimes V^{\otimes(k-j-2)}. \end{aligned}$$

Thus $e_i \cdot v_1 \wedge v_2 \wedge \cdots \wedge v_k = 0$ in the $U_{r,s}(\mathfrak{sl}_n)$ module $V^{\otimes k} / \sum_{i=0}^{k-2} V^{\otimes i} \otimes S_{r,s}^2(V) \otimes V^{\otimes(k-i-2)}$.

From the above discussion, we have that the fundamental representation $V(\overline{\omega}_k)$ is isomorphic to a submodule of $V^{\otimes k} / \sum_{i=0}^{k-2} V^{\otimes i} \otimes S_{r,s}^2(V) \otimes V^{\otimes(k-i-2)}$. By taking special values of r and s , one sees that two modules have the same dimension, so they are isomorphic. $\square$

Acknowledgments

NJ gratefully acknowledges the partial support of Simons Foundation grant 198129 and NSFC grant 11271138 during this work.

REFERENCES

- [1] G. Benkart and S. Witherspoon, *A Hopf structure for down-up algebras*, Math. Z. 238 (2001), 523–553.
- [2] G. Benkart and S. Witherspoon, *Two-parameter quantum groups and Drinfeld doubles*, Algebr. Represent. Theory 7 (2004), 261–286.
- [3] G. Benkart and S. Witherspoon, *Representations of two-parameter quantum groups and Schur-Weyl duality*, In: Hopf algebras, Lecture Notes in Pure and Appl. Math., 237, New York:Dekker, 2004, pp. 62–92.
- [4] J. Du, B. Parshall and J. Wang, *Two-parameter quantum linear groups and the hyperbolic invariance of q -Schur algebras*, J. London Math. Soc. (2)44 (1991), 420–436.
- [5] R. Dipper and S. Donkin, *Quantum GL_n* , Proc. London Math. Soc. (3) 63 (1991), no. 1, 165–211.
- [6] V. G. Drinfeld, *Quantum groups*, Proc. ICM, Vol. 1, 2 (Berkeley, Calif., 1986), pp.798–820, Amer. Math. Soc., Providence, RI, 1987.
- [7] M.-L. Ge, Y.-S. Wu and K. Xue, *Explicit Triognometric Yang-Baxterization*, Inter. J. of Modern Phys. A 6 (1991), 3735–3779.
- [8] N. Hu, M. Rosso and H. Zhang, *Two-parameter quantum affine algebra $U_{r,s}(\widehat{\mathfrak{sl}}_n)$, Drinfeld realization and quantum affine Lyndon basis*, Commun. Math. Phys. 278 (2008), 453–486.
- [9] M. Jimbo, *A q -difference analogue of $U(g)$ and the Yang-Baxter equation*, Lett. Math. Phys. 10 (1985), 63–69.
- [10] N. Jing, *Quantum groups with two parameters*, Contemp. Math. 134 (1992), 129–138.
- [11] N. Jing, K. Misra and M. Okado, *q -wedge modules for quantized enveloping algebras of classical type*, J. Algebra 230 (2000), 518–539.
- [12] S.-J. Kang, M. Kashiwara, K. C. Misra, T. Miwa, T. Nakashima, and A. Nakayashiki, *Perfect crystals of quantum affine Lie algebras*, Duke Math. J. 68 (1992), 499–607.
- [13] Y. Manin, *Topics in noncommutative geometry*, Princeton Univ. Press, 1991.
- [14] M. Rosso, *Finite dimensional representations of the quantum analog of the enveloping algebra of a complex simple Lie algebra*, Commun. Math. Phys. 117 (1988), 581–593.
- [15] M. Takeuchi, *A two-parameter quantization of $GL(n)$* , Proc. Japan. Acad. 66 Ser. A (1990), 112–114.

DEPARTMENT OF MATHEMATICS, NORTH CAROLINA STATE UNIVERSITY, RALEIGH, NC 27695, USA

SCHOOL OF SCIENCES, SOUTH CHINA UNIVERSITY OF TECHNOLOGY, GUANGZHOU 510640, CHINA

CHERN INSTITUTE OF MATHEMATICS, NANKAI UNIVERSITY, TIANJIN 300071, CHINA